

UNIT-4

①

Elementary combinatorics (permutation & combinations)

combinatorics :- combinatorics is an important part of discrete mathematics that solves counting problems without actually enumerating all possible cases.

combinatorics deals with counting the number of ways of arranging or choosing objects from a finite set according to certain specified rules.

In other words, combinatorics is concerned with problems of permutations and combinations.

permutations :- (arrangement - purpose)

An ordered arrangement of r elements of a set containing n distinct elements is an r -permutation of n elements and is denoted by $P(n, r)$ or nPr where $r \leq n$.

$$P(n, r) = n(n-1)(n-2) \dots (n-r+1)$$

$$\boxed{\begin{aligned} P(n, r) &= \frac{n!}{(n-r)!} \\ P(n, n) &= n! \end{aligned}}$$

combinations: An ordered selection of r elements of a set containing n distinct elements is called an r -combination of n elements and is denoted by nC_r $\binom{n}{r}$ nC_r $({}^nC_r)$

$$\therefore {}^nC_r = \frac{n!}{r!(n-r)!}$$

$${}^nC_n = 1$$

permutation: An arrangement of elements of a set is called a permutation of elements.

$${}^nP_r = \frac{n!}{(n-r)!}$$

problems:-

① How many 4 digit numbers can be formed by using the digits 2, 4, 6, 8 when repetition of digit is allowed.

sol:- Since repetition is allowed, each of the 4 places in a 4 digit number can be filled up in 4 ways by

(2)

the given 4 digits

<u>4 ways</u>	<u>4 ways</u>	<u>4 ways</u>	<u>4 ways</u>
Thousands	Hundreds	Tens	Unit

the required no. of 4 digit numbers
 $= n^r \Rightarrow 4^4 \Rightarrow 4 \times 4 \times 4 \times 4$
 $= 256$

Example problems on permutations

(1) number of permutations of n distinct objects (without duplication) :

The number of different arrangements (permutations) of n distinct objects, taken all at a time is

$$P(n, n) = \frac{n!}{(n-n)!} = \frac{n!}{0!} = n!$$

$$\boxed{P(n, n) = n!}$$

(2) number of permutations of r objects among n distinct objects :

Suppose we are given n distinctly
and wish to arrange r of the objects
denoted by

$$p(n, r) = \frac{n!}{(n-r)!}$$

$${}_n P_r = P(n, r)$$

(3) Number of permutations of n distinctly
objects (with duplication) :-

It is required to find the no. of
permutations that can be formed from a
collection of n objects of which n_1 are of
one type, n_2 are of a second type, ..., n_k are
of k^{th} type with $n_1 + n_2 + \dots + n_k = n$
then the no. of permutations of n objects
are

$$\frac{n!}{n_1! * n_2! * \dots * n_k!}$$

(4) circular permutation :- permutations in
a circle are ~~asked~~ called circular permu-
tation.

the tot no. of ways of arranging
the n persons in a circle $= (n-1)!$

Here 'n' is the no. of permutations

problems

(3)

① How many ways are there to sit 10 boys and 10 girls around a circular table?

Sol: - Here 10 Boys & 10 girls are sit around a circular table.

Tot no. of persons = $10 + 10 = 20$
 $n = 20$

The tot no. of ways of circular permutations are $(n-1)!$
 $= (20-1)! = 19! = 19 \times 18 \times \dots \times 1$

② How many ways are there 3 persons sit around a round table?

Sol: - No. of persons $n = 3$
The tot no. of ways of arranging a round table is
3 persons
 $(n-1)! = (3-1)! = 2! = 2$

③ How many different arrangements of letters in the word BOUGHT?

Sol: - The given word BOUGHT contains 6 letters that are distinct (without repetition)

The tot no. of arrangements of letters
in the word BOUGHT = $P(n, n) = \frac{n!}{n-n!}$
 $= \frac{7!}{0!}$
 $= 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1$
 $= 720$

(4) How many different strings of
length 4 can be formed using the
letters of the word PROBLEM?

sol: - The given word PROBLEM
has 7 letters $\therefore n=7$
The no. of different strings of
length 4 can be formed by using the
letters of the word PROBLEM =

$$P(n, r) = \frac{n!}{(n-r)!} \quad P(7, 4) = \frac{7!}{(7-4)!} = \frac{7!}{3!}$$

$$= \frac{7 \times 6 \times 5 \times 4 \times 3!}{3!}$$

$$= 840$$

(5) How many words of three distinct
letters can be formed from the letters
of the word PASCAL?

sol: - The given word PASCAL
contains 6 letters

$$\therefore n=6$$

$$P(n, r) = P(6, 3) = \frac{6!}{(6-3)!} = \frac{6!}{3!}$$

$$= \frac{6 \times 5 \times 4 \times 3!}{3!} = 120 \text{ ways.}$$

$\therefore$ the no. of ways 3 distinct letters can be formed by using the letters of the word ~~pas~~ PASCAL = 120 ways.

(6) Find the no. of permutations of the letters of the word ENGINEERING.

Sol:- The given word has 11 letters out of which

$$\begin{aligned} E &= 3 \text{ } (n_1) \\ N &= 3 \text{ } (n_2) \\ I &= 2 \text{ } (n_3) \\ G &= 2 \text{ } (n_4) \\ R &= 1 \text{ } (n_5) \end{aligned}$$

$\therefore$ total no. of permutations of the letters of the word ENGINEERING =

$$= \frac{n!}{n_1! \times n_2! \times n_3! \times n_4! \times n_5!}$$

$$= \frac{11!}{3! \times 3! \times 2! \times 2! \times 1!}$$

$$= \frac{11!}{6 \times 6 \times 4} \Rightarrow \frac{11 \times 10 \times 9 \times 8 \times 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1}{6 \times 6 \times 4} = 2,77,200$$

H.W

① Find permutations of

$$(a) \text{ SUCCESS} = \frac{7!}{3! \times 2! \times 1! \times 1!} = 420$$

$$(b) \text{ STRUCTURES} = \frac{10!}{2! \times 2! \times 2! \times 2! \times 1!} = 2,268$$

$$(c) \text{ MATHEMATICS} = \frac{11!}{2! \times 2! \times 2! \times 4! \times 1!} = 49,89,600$$

① prove that $(2n)! = 2^n n! \{1 \cdot 3 \cdot 5 \dots (2n-1)\}$

sol:-

$$\begin{aligned}(2n)! &= 2n(2n-1)(2n-2) \dots 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \\&= [2n(2n-2) \dots 4 \cdot 2] (2n-1)(2n-3) \dots 5 \cdot 3 \cdot 1 \\&= [2^n (n-1) \dots 2 \cdot 1] \\&= 2^n (n-1)(n-2) \dots 2 \cdot 1 [1 \cdot 3 \cdot 5 \dots (2n-1)] \\&= 2^n n! [1 \cdot 3 \cdot 5 \dots (2n-1)]\end{aligned}$$

② If $nP_2 = 72$ find the value of n .

sol:- Here $nP_2 = 72 \Rightarrow \frac{n!}{(n-2)!} = 72$

$$\frac{n(n-1)(n-2)!}{(n-2)!} = 72$$

$$n(n-1) = 72$$

$$n - n = 72 \Rightarrow n - n - 72 = 0$$

$$n^2 - n - 72 = 0$$

$$n(n-9) + 8(n-9) = 0$$

$$(n+8)(n-9) = 0$$

$$n = -8, n = 9$$

But $n \neq -8$ (n cannot be negative)

$$\therefore \boxed{n=9}$$

② $\therefore {}^{2n+1}P_{n-1} : {}^{2n-1}P_n = 3/5$ find n ;

③ $\therefore {}^{2n+1}P_{n-1} = \frac{(2n+1)!}{(n+2)!}$ and ${}^{2n-1}P_n = \frac{(2n-1)!}{(n-2)!}$

$$\therefore {}^{2n+1}P_{n-1} \div {}^{2n-1}P_n = \frac{(2n+1)!}{(n+2)!} \cdot \frac{(n-2)!}{(2n-1)!} = \frac{3}{5}$$

$$\frac{(2n+1)!}{(2n+1-n+1)!} = \frac{3}{5}$$

$$\frac{(2n+1)!}{(2n-1)!} = \frac{3}{5}$$

$$\frac{(2n+1)!}{(n+2)!} \cdot \frac{(n-1)!}{(2n-1)!} = \frac{3}{5}$$

$$\frac{(2n+1)(2n+1-1)(2n+1-2)!}{(n+2)(n+2-1)(n+2-2)(n+2-3)!} \cdot \frac{(n-1)!}{(2n-1)!} = \frac{3}{5}$$

$$\frac{(2n+1)(2n)(2n-1)!}{(n+2)(n+1)(n)(n-1)!} \cdot \frac{(n-1)!}{(2n-1)!} = \frac{3}{5}$$

$$\frac{(4n+2)}{(n^2+3n+2)} = \frac{3}{5}$$

$$\begin{aligned} \Rightarrow 20n+10 &= 3n^2+6n+6 \\ \Rightarrow 3n^2+9n-20n+6-10 &= 0 \\ \Rightarrow 3n^2-11n-4 &= 0 \end{aligned}$$

$$(n-4)(3n+1) = 0$$

$$\therefore n = 4, \text{ or } -\frac{1}{3}$$

~~3-4-4~~

$$\text{But } n \neq -\frac{1}{3} \therefore n = 4$$

(3) In how many ways can the letters of the word DISCRETE so that the vowels may come together?

sol: - no. of letters in the given word $n = 8$
 vowels in the word $= 3$

$$\therefore \text{The required number} = 3! \times (8-3)!$$

$$= 3! \times 5!$$

$$= 720$$

(4) In how many ways can 7 boys and 5 girls be seated in a row so that no two girls sit together.

sol: - since there is no restriction on boys, first of all we fix the position of 7 boys.

Their positions are indicated as

$$\times B_1 \times B_2 \times B_3 \times B_4 \times B_5 \times B_6 \times B_7 \times$$

7 boys can be arranged in $7!$ ways.

now if 5 girls sit at places
including the two ends indicated by X,
then no two of the 5 girls will
sit together. ⑥

clearly, 5 girls can be seated on
8 places in 8P_5 ways.

Hence the required no. of ways of
seating 7 boys and 5 girls under the
given condition

$$= {}^8P_5 \times 7!$$

$$= \frac{8!}{3!} \times 7!$$

$$= 8! \times \frac{7 \times 6 \times 5 \times 4 \times 3!}{3!}$$

$$= 40320 \times 24$$

$$= 96768$$

combinations

(with repetition) ⑦

Suppose we wish to select a combination of r objects with repetition from a set of n distinct objects. The no. of such selections is given by

$$C(n+r-1, r) = C(r+n-1, n-1)$$

→ The following are the other interpretations of this

(i) $C(n+r-1, r) = C(r+n-1, n-1)$ represents the no. of ways in which r identical objects can be distributed among n distinct containers.

(ii) $C(n+r-1, r) = C(r+n-1, n-1)$ represents the no. of non-negative integer solutions of the equation

Eg:- In how many ways can 20 similar books be placed on 5 diff shelves?

Here $r=20, n=5$

Req no. of way $\Rightarrow$

$$\begin{aligned} &= C(n+r-1, r) \\ &= C(5+20-1, 20) \\ &= C(24, 20) \\ &= 101626 \end{aligned}$$

n - distinct elements
 $\downarrow$
 r - identical object w.r
 $\downarrow$
no. of such selections
 $C(n+r-1, r) = C(r+n-1, n-1)$
 $\swarrow \searrow$
 n distinct identical objects no. of non-negative integers

Note :- A non-negative integer solution of the equation $x_1 + x_2 + \dots + x_n = r$ is an n -tuple, where $x_1, x_2, x_3, \dots, x_n$ are non-negative integers whose sum is r .

eg:- ① In how many ways can we distribute 10 identical marbles among 6 distinct containers?

sol:- comparing with st interpretation

10 identical marbles $r = 10$
 6 distinct containers $n = 6$
 no. of such selections

$$c(r+n-1, r) = c(10+6-1, 10)$$

$$= c(15, 10)$$

$$= \frac{15!}{10! 5!}$$

$$= 3003$$

eg② :- Find the no. of non-negative integer solutions of the equation

sol:- sum of non-negative integers is 8
 i.e. $x_1 + x_2 + x_3 + x_4 + x_5 = 8$
 No. of non-negative integers $n=5$
 $r=8$

$$c(n+r-1, r) = c(5+8-1, 8)$$

$$= c(12, 8)$$

$$= \frac{12!}{8! 4!} = 495$$

③ In how many ways can we distribute 12 identical pencils to 5 children so that every child gets at least one pencil?

sol :- No. of pencils = 12
 No. of children $n = 5$
 every child get at least one pencil
 means each child may get ≥ 1 pencil
 First, we distribute one pencil to each child, then the remaining identical pencils $(12 - 5 = 7)$ can be distributed to 5 children.

$\therefore$ no. of ways to distribute 7 identical pencils to 5 children =

$$r = 7, n = 5$$

$$C(r+n-1, r) = C(7+5-1, 7)$$

$$= C(11, 7)$$

$$= \frac{11!}{7!4!}$$

$$= 330 \text{ ways}$$

④ In how many ways can we distribute 7 apples and 6 oranges among 4 children so that each child gets one apple?

sol :- No. of apples = 7

No. of oranges = 6

No. of children = 4

each children gets at least one apple,
i.e. every apple may get ≥ 1 apple.

first we distribute one apple to
each child. The remaining apples
($6-4=2$) 2 can be distributed to 4
children.

No. of ways to distribute 2 apples
to 4 children

$$= {}^nC_{(r+(n-1), n)} \quad \begin{matrix} r=2 \\ n=4 \end{matrix}$$

$$= {}^nC_{(6, 2)}$$

$$= \frac{6!}{2! 4!}$$

$$= 15 \text{ ways.}$$

No. of ways to distribute 6
oranges to 4 children

$$= {}^nC_{(r+n-1, r)} \quad \begin{matrix} r=6 \\ n=4 \end{matrix}$$

$$= {}^nC_{(6+4-1, 6)}$$

$$= {}^nC_{(9, 6)}$$

$$= \frac{9!}{6! 3!}$$

$$= 84 \text{ ways}$$

$\therefore$ Total no. of ways to distribute
the fruits under given condition 84 ways

→ find the no. of positive integer solutions of the equation (9)
 $x_1 + x_2 + x_3 = 17$ where $x_1 \geq 1, x_2 \geq 1, x_3 \geq 1$

Sol:- Given equation $x_1 + x_2 + x_3 = 17$

where $x_1 \geq 1, x_2 \geq 1, x_3 \geq 1$

let us consider three non-negative integers

y_1, y_2 and y_3

$$y_1 = x_1 - 1 \Rightarrow x_1 = y_1 + 1$$

$$y_2 = x_2 - 1 \Rightarrow x_2 = y_2 + 1$$

$$y_3 = x_3 - 1 \Rightarrow x_3 = y_3 + 1$$

the above eqn can be defined in terms of y_1, y_2, y_3

$$x_1 + x_2 + x_3 = 17$$

$$(y_1 + 1) + (y_2 + 1) + (y_3 + 1) = 17$$

$$y_1 + y_2 + y_3 + 3 = 17$$

$$y_1 + y_2 + y_3 = 14$$

$$\therefore y_1 + y_2 + y_3 = 14$$

no. of positive integer sol^{ns} of the above equation = $C(n+r-1, r)$

$$= C(3+14-1, 14)$$

$$= C(16, 14)$$

$$= \frac{16!}{14! 2!}$$

$$= 120$$

$\therefore$ no. of positive integer sol^{ns} of the given eqⁿ = 120

e.

$$\begin{matrix} n=3 \\ r=14 \end{matrix}$$

H.W

-> Find the no. of integer solⁿ of
the eqⁿ $x_1 + x_2 + x_3 + x_4 + x_5 = 30$
where $x_1 \geq 2, x_2 \geq 3, x_3 \geq 4, x_4 \geq 2, x_5 \geq 0$

Sol:- Given equation $x_1 + x_2 + x_3 + x_4 + x_5 = 30$
let us consider the +ve integers

y_1, y_2, y_3, y_4 & y_5

Now let us set-

$$y_1 = x_1 - 2 \Rightarrow x_1 = y_1 + 2$$

$$y_2 = x_2 - 3 \Rightarrow x_2 = y_2 + 3$$

$$y_3 = x_3 - 4 \Rightarrow x_3 = y_3 + 4$$

$$y_4 = x_4 - 2 \Rightarrow x_4 = y_4 + 2$$

$$y_5 = x_5 \Rightarrow x_5 = y_5$$

the above eqn can be written as

$$x_1 + x_2 + x_3 + x_4 + x_5 = 30$$

$$(y_1 + 2) + (y_2 + 3) + (y_3 + 4) + (y_4 + 2) + y_5 = 30$$

$$y_1 + y_2 + y_3 + y_4 + y_5 = 30 - 11 = 19$$

$$\therefore y_1 + y_2 + y_3 + y_4 + y_5 = 19$$

the no. of integer solⁿ of the above

$$\text{eqⁿ } = C(n+r-1, r)$$

$$= C(5+19-1, 19)$$

$$= C(23, 19)$$

$$= \frac{23!}{19! 4!} = 8855$$

$$\begin{matrix} n=5 \\ r=19 \end{matrix}$$

properties

(10) (10)

$$(1) \quad nC_r = nC_{n-r} \quad (0 \leq r \leq n)$$

pf:- we have $nC_{n-r} = \frac{n!}{(n-r)! (n-n+r)!}$

$$= \frac{n!}{(n-r)! r!} = nC_r$$

$$(2) \quad nC_r + nC_{r-1} = n+1C_r \quad (0 \leq r \leq n)$$

sw:- $nC_r + nC_{r-1} = \frac{n!}{r! (n-r)!} + \frac{n!}{(r-1)! (n-r+1)!}$

$$= n! \left[\frac{1}{r! (n-r)!} + \frac{1}{(r-1)! (n-r+1)!} \right]$$

$$= n! \left[\frac{1}{(n-r)! r (r-1)!} + \frac{1}{(n-r+1) (n-r+1-1)! (r-1)!} \right]$$

$$= n! \left[\frac{1}{r (r-1)! (n-r)!} + \frac{1}{(n-r+1) (n-r)! (r-1)!} \right]$$

$$= \frac{n!}{(n-r)! (r-1)!} \left[\frac{1}{r} + \frac{1}{n-r+1} \right]$$

$$= \frac{n!}{(n-r)! (r-1)!} \left[\frac{n-r+1 + r}{r (n-r+1)} \right]$$

$$= \frac{n!}{(n-r)! (r-1)!} \left[\frac{(n+1)}{r (n-r+1)} \right]$$

$$= \frac{(n+1) n!}{x(x-1)! (n-x+1) (n-x)!}$$

$$= \frac{(n+1)!}{x! (n-x+1)!}$$

$$nC_x + nC_{x-1} = n+1 C_x$$

$\Rightarrow \dots$

③ $nC_x = nC_y \Rightarrow x=y$ or $x+y=n$

say:- we have $nC_x = nC_y$

$$\Rightarrow nC_x = nC_{n-y} \quad \left[\because nC_x = nC_{n-x} \right]$$

$$\Rightarrow x=y \text{ (or) } x=n-y$$

$$\Rightarrow x=y \text{ (or) } x+y=n.$$

Relation b/w nC_x and $n+1 C_x$

$$\frac{n+1 C_x}{nC_x} = \frac{\frac{(n+1)!}{x! (n+1-x)!}}{\frac{n!}{x! (n-x)!}} = \frac{(n+1)!}{x! (n+1-x)!} \times \frac{x! (n-x)!}{n!}$$

$$= \frac{(n+1) (n+1-1)!}{(n+1-x) (n+1-x-1)!} \times \frac{(n-x)!}{n!}$$

$$= \frac{n+1}{(n+1-x)}$$

Q Evaluate the following

(119)

$$(i) {}^6C_3 = \frac{6!}{3!(6-3)!} = \frac{6!}{3!3!} = \frac{6 \times 5 \times 4 \times 3 \times 2 \times 1}{3 \times 2 \times 1 \times 3 \times 2 \times 1} = 20$$

$$(ii) {}^{10}C_8 = {}^{10}C_{10-8} = {}^{10}C_2 = \frac{10!}{2!8!} = 45$$

(2) compute 8P_5 & 6C_3

$$\text{sol: } {}^8P_5 = \frac{8!}{(8-5)!} = \frac{8!}{3!} = \frac{8 \times 7 \times 6 \times 5 \times 4 \times 3!}{3!} = 6720$$

$${}^6C_3 = \frac{6!}{3!(6-3)!} = \frac{6!}{3!3!} = 20$$

(3) ${}^{18}C_r = {}^{18}_{r+2}$ find the value of nC_5

$$\text{sol: } \therefore {}^nC_r = {}^nC_{n-r} \Rightarrow r + n - r = n$$

$$\text{so } r + r + 2 = 18 \Rightarrow 2r + 2 = 18$$

$$\Rightarrow r + 1 = 9$$

$$r = 8$$

$$\therefore {}^nC_5 = {}^8C_5 = \frac{8!}{5!(8-5)!} = \frac{8!}{5!3!} = \frac{8 \times 7 \times 6 \times 5!}{5! \times 3 \times 2 \times 1} = 56$$

H.W

pg-429

→ Find the no. of integer solutions of $x_1 + x_2 + x_3 + x_4 + x_5 = 30$
where $x_1 \geq 2, x_2 \geq 3, x_3 \geq 4, x_4 \geq 2, x_5 \geq 0$

(4) If $nC_n = 56$ & $nP_n = 336$ find n

sol :- $nC_n = 56 \Rightarrow \frac{n!}{n!(n-n)!} = 56$

$nP_n = 336 \Rightarrow \frac{n!}{(n-n)!} = 336$

$\frac{nC_n}{nP_n} = \frac{56}{336} \Rightarrow \frac{n!}{n!(n-n)!} \times \frac{(n-n)!}{n!} = \frac{56}{336} = \frac{1}{6}$

$= \frac{1}{n!} = \frac{1}{6}$

$n! = 3!$

$\boxed{n = 3}$

Agueen $nP_n = 336$

$nP_3 = 336 \Rightarrow \frac{n!}{(n-3)!} = 336$

$\frac{n(n-1)(n-2)(\cancel{n-3})!}{(\cancel{n-3})!} = 336$

$n(n-1)(n-2) = 8 \times 7 \times 6$

$\therefore \boxed{n = 8}$

(5) If $1000C_{98} = 999C_{97} + nC_{901}$; find n

sol :- Here $1000C_{98} = 999C_{97} + nC_{901}$

(31) $1000C_{902} = 999C_{902} + nC_{901} \quad [\because nC_n = nC_{n-n}]$

(32) $999+1C_{902} = 999C_{902} + nC_{902-1}$

$${}^{n+1}C_n = {}^nC_n + {}^nC_{n-1}$$

$$\therefore n = 999$$

6) In how many ways can 4 questions be selected from 7 questions?

Sol :- The required no. of ways

$${}^7C_4 = \frac{7!}{4!(7-4)!} = \frac{7!}{4!3!} = 35$$

7) In how many ways can you select at least one king, if you choose five cards from a deck of 52 cards?

Sol :- There are 4 kings in a deck of 52 cards

to no. of ways of choosing	1 king =	${}^4C_1 \times {}^{48}C_4$
" " " " " "	2 " =	${}^4C_2 \times {}^{48}C_3$
" " " " " "	3 " =	${}^4C_3 \times {}^{48}C_2$
" " " " " "	4 " =	${}^4C_4 \times {}^{48}C_1$

$$\therefore \text{The required number } {}^4C_1 \times {}^{48}C_4 + {}^4C_2 \times {}^{48}C_3 + {}^4C_3 \times {}^{48}C_2 + {}^4C_4 \times {}^{48}C_1$$

$$= 886, 656$$

Note: Total no. of ways in which one or more things are taken $2^n - 1$

eg:- ① you have 4 friends, in how many ways can you invite one or more them for dinner?

say:- you may invite 1, 2, 3, or 4 of your friends to dinner. Hence the required no. of ways

$$= 2^n - 1 \\ = 2^4 - 1 \\ = 63.$$

② There are 5 questions in a question paper. In how many ways can a boy solve one or more questions?

say:- $2^5 - 1 = 31$

Note:- the selection made of things where p, q, r or of third kind and so on, the tot. no. of possible selections is

$$[(p+1)(q+1)(r+1)\dots] - 1$$

eg:- In how many ways can a selection be made out of 3 mangoes, 5 oranges and 2 apples?

$$p = 3, q = 5, r = 2$$

say:- Here $\therefore$ Req. no. of combinations

$$(3+1)(5+1)(2+1) - 1 \\ = 4 \cdot 6 \cdot 3 - 1 \\ = 72 - 1 = 71$$

pigeonhole principle

(13)

pigeonhole principle : — If n pigeons are accommodated in m pigeonholes and $n > m$ then at least one pigeonhole will contain two or more pigeons.

Generalized pigeonhole principle :

If n pigeons are accommodated in m pigeonholes and $n > m$ then one of the pigeon holes must contain at least $\left\lfloor \frac{(n-1)}{m} \right\rfloor + 1$ pigeons.

eg:- ① P.T in any set of 29 persons at least 5 persons must have been born on the same day of the week.

sol:- A week contains 7 days : 7 pigeon holes
a set of 29 persons : 29 pigeons

According to generalized pigeonhole principle

$$\left(\frac{n-1}{m} \right) + 1 \Rightarrow \frac{29-1}{7}$$

$$n = 29$$

$$m = 7$$

$$= \frac{28}{7} = 4$$

Among a set of 29 persons, 5 persons must have been born on the same day of the week

(2) Find the minimum no. of 12 students in a class to be sure that four out of them are born ~~on~~ ⁱⁿ the same day of a week.

Sol: - we consider each month as a pigeonhole. then $m = 12$
 we have to find the minimum no. of students (pigeons) so that four out of them are born in the same month.

$$\left\lceil \frac{n-1}{m} \right\rceil + 1 = 4$$

$$\frac{n-1}{m} = 3 \quad \Rightarrow \quad 3m = n-1$$

$$3(12) = n-1$$

$$36+1 = n \Rightarrow n = 37$$

$\therefore$ which is req. min no. of students

(3) If 9 books are to be kept in 4 shelves, there must be atleast one shelf which contains at least 3 books.

Sol: - $n = 9, m = 4$

By pigeon hole principle $\left\lceil \frac{n-1}{m} \right\rceil + 1 = 3$

$$\begin{aligned} \left(\frac{9-1}{4} \right) + 1 &= 3 \\ \frac{8}{4} + 1 &= 3 \\ 2 + 1 &= 3 \end{aligned}$$

~~there~~ so at least 3 of the persons have
 atleast one shelf will contain
 atleast 3 books
 =

(14)

principle of Inclusion - Exclusion

If A & B are any two finite sets

Then $|A \cup B| = |A| + |B| - |A \cap B|$

where $|A|$ denotes the cardinality of A .

$A = \{a, b, c\}$
 $|A| = 3$
 cardinality of A

If A, B, C are any 3 sets,

$$|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |B \cap C| - |A \cap C| + |A \cap B \cap C|$$

If A and B are two disjoint sets

then $|A \cup B| = |A| + |B|$

for generalisation If $A_1, A_2, \dots, A_n$ are finite sets then

$$|A_1 \cup A_2 \cup \dots \cup A_n| = \sum_{1 \leq i \leq n} |A_i| - \sum_{1 \leq i < j \leq n} |A_i \cap A_j| + \sum_{1 \leq i < j < k \leq n} |A_i \cap A_j \cap A_k| - \dots + (-1)^{n+1} |A_1 \cap A_2 \cap \dots \cap A_n|$$

no. of elements

Give a formula for the union of four sets.

eg- ① on the A_1, A_2, A_3, A_4

$$\therefore |A_1 \cup A_2 \cup A_3 \cup A_4| = |A_1| + |A_2| + |A_3| + |A_4| -$$

$$\begin{aligned}
 & - |A_1 \cap A_2| - |A_1 \cap A_3| - |A_1 \cap A_4| - |A_2 \cap A_3| - |A_2 \cap A_4| - \\
 & |A_3 \cap A_4| + |A_1 \cap A_2 \cap A_3| + |A_1 \cap A_2 \cap A_4| + \\
 & |A_1 \cap A_3 \cap A_4| + |A_2 \cap A_3 \cap A_4| - \\
 & |A_1 \cap A_2 \cap A_3 \cap A_4|
 \end{aligned}$$

problems

(i) In a sample of 200 logic chips, 46 have a defect D_1 , 52 have a defect D_2 , 60 have defect D_3 , 14 have defects D_1 and D_2 , 16 have defects D_2 and D_3 , and 3 have all the three defects. Find the no. of chips having
 (i) at least one defect
 (ii) no defect

say:- let U denotes the set of all chips in a given sample, and A, B, C denote the set of chips having defects D_1, D_2, D_3 respectively.

Then we have

$$\begin{aligned}
 |U| &= 200, & |A| &= 46, & |B| &= 52, & |C| &= 60, & |A \cap B| &= 14 \\
 |A \cap C| &= 16, & |B \cap C| &= 20 & \text{and} & |A \cap B \cap C| &= 3
 \end{aligned}$$

(i) The set of chips having at least one defect is $A \cup B \cup C$

∴ the no. of chips having at least one defect = $|A \cup B \cup C|$

$$\begin{aligned} &= |A| + |B| + |C| - |A \cap B| - |B \cap C| - |C \cap A| + |A \cap B \cap C| \\ &= 46 + 52 + 60 - 14 - 16 - 20 + 3 \\ &= 161 - 50 \\ &= 111 \end{aligned}$$

② write the principle of Inclusion-Exclusion from a group of 10 professors. How many ways can committees of 5 members be formed so that at least one professor A and professor B will be included.

Sol: - The total no. of committees $C(10, 5)$ let A & B be the sets of committees that include professor A and professor B respectively

$$\begin{aligned} \text{Then } |A| &= C(9, 4) \text{ \& } |B| = C(9, 4) \text{ \& } \\ |A \cap B| &= C(8, 3) \end{aligned}$$

By the principle of the inclusion-exclusion

$$\begin{aligned} |A \cup B| &= |A| + |B| - |A \cap B| \\ &= C(9, 4) + C(9, 4) - C(8, 3) \\ &= 2C(9, 4) - C(8, 3) \end{aligned}$$

$$= 2 \times \frac{9!}{4!5!} - \frac{8!}{3!5!}$$

$$= 2 \times \frac{9 \times 8 \times 7 \times 6}{24} - \frac{8 \times 7 \times 6}{6}$$

$$= 252 - 56$$

$$= 196.$$

③ How many integers between 1 & 300 (inclusive) are divisible by at least one of 5, 6 and 8.

sol: - Let $S = \{1, 2, 3, \dots, 300\}$ then

$|S| = 300$
Let A, B, C be the subset of S whose elements are divisible by 5, 6, 8 respectively.
Then we have,

$$|A| = \left\lfloor \frac{300}{5} \right\rfloor = 60, \quad |B| = \left\lfloor \frac{300}{6} \right\rfloor = 50$$

$$|C| = \left\lfloor \frac{300}{8} \right\rfloor = 37, \quad |A \cap B| = \left\lfloor \frac{300}{30} \right\rfloor = 10$$

($\because$ LCM of 5 & 6 is 30)

$$|A \cap C| = \left\lfloor \frac{300}{40} \right\rfloor = 7, \quad |B \cap C| = \left\lfloor \frac{300}{24} \right\rfloor = 12$$

($\because$ LCM of 6 & 8 is 24)

$$\text{and } |A \cap B \cap C| = \left\lfloor \frac{300}{120} \right\rfloor = 2 \quad (\because \text{LCM of 5, 6, 8 is 120})$$

By the principle of inclusion exclusion ⁽¹⁶⁾

we have

$$\begin{aligned} |A \cup B \cup C| &= |A| + |B| + |C| - |A \cap B| - |B \cap C| \\ &\quad - |C \cap A| + |A \cap B \cap C| \\ &= 60 + 50 + 37 - 10 - 7 - 12 + 2 \\ &= 120 \end{aligned}$$

Thus 120 elements of S are divisible
by at least one of 5, 6, 8.

BINOMIAL THEOREM

A Binomial theorem describes the algebraic expansion of powers of a Binomial with two variables.

$$(x+y)^n = nC_0 x^n y^0 + nC_1 x^{n-1} y^1 + nC_2 x^{n-2} y^2 + \dots \\ \dots + nC_{n-1} x^1 y^{n-1} + nC_n x^0 y^n$$

It can be written as

$$(x+y)^n = \sum_{r=0}^n nC_r x^{n-r} y^r$$

$$\rightarrow (1+x)^n = nC_0 1^n x^0 + nC_1 1^{n-1} x^1 + nC_2 1^{n-2} x^2 + \dots \\ nC_3 1^{n-3} x^3 + \dots$$

$$=) nC_0 x^n + nC_1 x^{n-1} + nC_2 x^{n-2} + nC_3 x^{n-3} + \dots$$

$$\Rightarrow 1 \cdot x^n + \frac{n!}{1!(n-1)!} x^{n-1} + \frac{n!}{2!(n-2)!} x^{n-2} + \frac{n!}{3!(n-3)!} x^{n-3} + \dots$$

$$=) 1 + \frac{n(n-1)!}{(n-1)!} x + \frac{n(n-1)(n-2)!}{2!(n-2)!} x^2 + \dots$$

$$\frac{n(n-1)(n-2)(n-3)!}{3!(n-3)!} x^3 + \dots$$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!} x^2 + \frac{n(n-1)(n-2)}{3!} x^3 + \dots$$

problem :

① find out the coefficient of $x^9 y^3$ in $(x+2y)^{12}$

sol : — we know Binomial Theorem
 $(x+y)^n = \sum_{r=0}^n nC_r \cdot x^{n-r} \cdot y^r$
 according to the Binomial theorem

$$x = x, \quad y = 2y, \quad n = 12$$

$$(x+2y)^{12} = \sum_{r=0}^{12} {}^{12}C_r x^{12-r} (2y)^r$$

$$= \sum_{r=0}^{\infty} {}^{12}C_r x^{12-r} 2^r y^r \quad \text{--- (1)}$$

we have to find out $x^9 y^3$

$$x^9 = x^{12-3} \quad / \quad y^n = y^{12-3} y^3$$

$$9 = 12 - 3$$

$$n = 3$$

$$n = 3$$

x value substitute in eq (1)

$${}^{12}C_3 \cdot 2^3 \cdot x^{12-3} y^3$$

$${}^{12}C_3 \cdot 2^3 \cdot x^9 y^3$$

$$\frac{12!}{3!9!} \times 2^3$$

$$\frac{12 \times 11 \times 10 \times 9!}{3 \times 6 \times 9!}$$

$$= 1760$$

$\therefore$ coefficient of $x^9 y^3$ in $(x+2y)^{12}$

$$= 1760$$

(2) Find out the coefficient of $x^5 y^2$ in the expansion of $(2x-3y)^7$

we know that, from Binomial theorem

$$(x+y)^n = \sum_{r=0}^{\infty} {}^nC_r x^{n-r} y^r$$

$$x = 2x, \quad y = -3y, \quad n = 7$$

$$(2x-3y)^7 = \sum_{r=0}^{\infty} {}^7C_r (2x)^{7-r} (-3y)^r$$

$$= \sum_{r=0}^{\infty} {}^7C_r 2^{7-r} x^{7-r} (-3)^r y^r \quad \text{--- (1)}$$

we have to find out the coefficient of

$$\begin{array}{l|l} n^5 \cdot y^2 & y^r = y^2 \\ n^{7-r} = n^5 & r = 2 \\ 7-r = 5 & \\ r = 2 & \\ r = 2 & \end{array}$$

$\therefore$ r value substitute in eqⁿ ①

$${}^7C_2 (2)^{7-2} (-3)^2 x^{7-2} y^2$$

$$= {}^7C_2 2^5 \cdot 9 \cdot x^5 y^2$$

$$\frac{7!}{2!5!} 32 \cdot 9 x^5 y^2$$

$$\frac{7 \times 6 \times 5!}{2 \times 1 \times 5!} 32 \cdot 9 x^5 y^2$$

$$= 21 \times 32 \times 9 \cdot x^5 y^2$$

$$= 6048 x^5 y^2$$

$\therefore$ coefficient of $x^5 y^2$ in $(2x-3y)^7$ is 6048

③ what is the coefficient of $x^2 y^4$ in $(x+y)^6$?

$$(x+y)^n = \sum_{r=0}^n {}^nC_r x^{n-r} y^r, \quad n=6.$$

$$(x+y)^6 = \sum_{r=0}^6 {}^6C_r x^{6-r} y^r \rightarrow \text{①}$$

we have to find out the coefficient of $x^2 y^4$

$$n^{6-r} = n^2$$

$$6-r = 2$$

$$r = 6-2 = 4$$

$$y^r = y^4$$

$$r = 4$$

substitute this value in eqn (1) (18)

$$6C_4 x^{6-4} y^4$$

$$6C_4 x^2 y^4$$

$$\frac{6!}{4!2!} x^2 y^4 = \frac{6 \times 5 \times 4!}{4! \times 2!} = 15$$

(9) x^{101} y^{99} in the expansion $(2x-3y)^{200}$?

$$(x+y)^n = \sum_{r=0}^{\infty} nC_r x^{n-r} y^r$$

$$n=2x, y=-3y, n=200$$

$$(2x-3y)^{200} = \sum_{r=0}^{\infty} 200C_r (2x)^{n-r} (-3y)^r \rightarrow (1)$$

$$= \sum_{r=0}^{\infty} 200C_r \frac{2^{n-r}}{2} (-3)^r y^r$$

$$x^{101} = x^{200-r}$$

$$y^r = y^{99}$$

$$200-r=101$$

$$r=99$$

$$200-101=99$$

$$r=99$$

$$= {}^{200}C_{99} \frac{2^{200-99}}{2} x^{200-99} (-3)^{99} (y)^{99}$$

$$= {}^{200}C_{99}$$

$$\frac{2^{101}}{2} 3^{99} \cdot y^{99} x^{101}$$

$$- {}^{200}C_{99}$$

$$x^{101} y^{99} \therefore - {}^{200}C_{99} \frac{2^{101}}{2} 3^{99}$$

coefficient of

=

Multinomial Theorem

Multinomial theorem is a generalization of the Binomial theorem with more than two variables.

$$(x_1 + x_2 + \dots + x_m)^n = \sum_{n_1, n_2, \dots, n_k} \frac{n!}{n_1! \times n_2! \times \dots \times n_k!}$$

where $n_1 + n_2 + \dots + n_k = n$

eg:- ① compute the following

(a) $\binom{7}{2, 3, 2}$

It is in the form of

$\binom{n}{n_1, n_2, n_3}$ where $n_1 + n_2 + n_3 = n$

By applying the multinomial theorem

$$\frac{n!}{n_1! \times n_2! \times n_3!} \Rightarrow \frac{7!}{2! \times 3! \times 2!} = 210$$

(b) $\binom{4}{1, 1, 2}$

this is in the form of

$\binom{n}{n_1, n_2, n_3}$

$$\frac{4!}{1! \times 1! \times 2!} = \frac{24}{2} = 12$$

(c) $\binom{12}{5, 3, 2, 2}$ It is in the form of $\binom{n}{n_1, n_2, n_3, n_4}$

By multinomial theorem $\frac{12!}{5! \times 3! \times 2! \times 2!} = 166320$

Q. what is the coefficient of $x^3y^2z^2$ in $(x+y+z)^9$?

say: - Buy the multinomial theorem $\left(\frac{n!}{n_1! \times n_2! \times n_3!} \right)$

$$\frac{9!}{3! \times 2! \times 2!} = 15120.$$

⑤ Determine the coefficient of xy^2z^2 in the expansion of $(x+y+z)^4$ using the theorem

expansion

say! - Buy Applying Meletano melle / Theorem

$$(n_1 + n_2 + n_3 + \dots + n_k)^n = \frac{n!}{n_1! \times n_2! \times \dots \times n_k!} (x_1)^{n_1} (x_2)^{n_2} \dots (x_k)^{n_k}$$

where $n_1 + n_2 + \dots + n_k = n$

hier $n_1 + n_2 + \dots$
 gegeben $n_2 = -3$, $(2n - 3 - 2)^4$

$n = 4, \quad x_1 = -2, \quad x_2 = -4, \quad x_3 = -2$

Apply multinomial theorem

$$\frac{n!}{n_1! \times n_2! \times n_3!} \cdot (x_1)^{n_1} (x_2)^{n_2} (x_3)^{n_3} = \frac{4!}{n_1! \times n_2! \times n_3!} (2x_1)^{n_1} (-4)^{n_2} (-z)^{n_3} \rightarrow (1)$$

we have to find out the coefficient of $x^{n_1} x^{n_2} x^{n_3}$

$$xyt^2 = x_1^{n_1} x_2^{n_2} x_3^{n_3}$$

$$\begin{aligned} n_1 &= 1 \\ n_2 &= 1 \\ n_3 &= 2 \end{aligned}$$

substitute n_1, n_2, n_3 value in eqn (1)

$$\begin{aligned} & \frac{4!}{1! \times 1! \times 2!} (2x)^1 (-y)^1 (-z)^2 \\ &= \frac{24}{1 \times 1 \times 2} \times 2^1 x^1 (-1)^1 y^1 (-1)^2 z^2 \\ &= 12 \times 2 \cdot x \cdot -y \cdot z^2 \\ &= -24 xy z^2 \\ \therefore \text{coefficient of } xy z^2 & \text{ is } -24 \end{aligned}$$

(2) determine the coefficient of $x^3 y^3 z^2$ in the expansion of $(2x - 3y + 5z)^8$
 sol:- By multinomial Theorem


$$\frac{n!}{n_1! \times n_2! \times n_3!} x_1^{n_1} x_2^{n_2} x_3^{n_3}$$

compare $n=8, \quad n_1=3 \quad x_1=2x$
 $n_2=3 \quad x_2=-3y$
 $n_3=2 \quad x_3=5z$

$$\begin{aligned} \therefore & \frac{8!}{3! \times 3! \times 2!} (2x)^3 (-3y)^3 (5z)^2 \\ &= \frac{8!}{3! \times 3! \times 2!} 2^3 (-3)^3 5^2 x^3 y^3 z^2 \\ &= -3024000 x^3 y^3 z^2 \\ \therefore \text{coefficient of } x^3 y^3 z^2 & \text{ is } -3024000 \end{aligned}$$

Pascal's Triangle

①



②

$$\begin{aligned}
 (a+b)^0 &= 1 \\
 (a+b)^1 &= 1a + 1b \\
 (a+b)^2 &= 1a^2 + 2ab + 1a^0b^2 \\
 (a+b)^3 &= 1a^3 + 3a^2b + 3ab^2 + 1a^0b^3 \\
 (a+b)^4 &= 1a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + 1a^0b^4
 \end{aligned}$$

Pascal's Identity for $0 < r < n$

$$\binom{n}{r} + \binom{n}{r-1} = \binom{n+1}{r}$$

$$\binom{n}{r} = \frac{n!}{r! (n-r)!}$$

$$\frac{n!}{r! (n-r)!} + \frac{n!}{(r-1)! (n-r+1)!}$$

$$\frac{n!}{r (r-1)! (n-r)!} + \frac{n!}{(r-1)! (n-r+1) (n-r+1-1)!}$$

$$\frac{n!}{r (r-1)! (n-r)!} + \frac{n!}{(r-1)! (n-r+1) (n-r)!}$$

$$\frac{n!}{(r-1)! (n-r)!} \left[\frac{1}{r} + \frac{1}{n-r+1} \right]$$

$$\frac{n!}{(r-1)! (n-r)!} \left[\frac{n-r+1 + r}{r (n-r+1)} \right]$$

$$\frac{n!}{(r-1)! (n-r)!} \left[\frac{n+1}{r (n-r+1)} \right]$$

$$\frac{n! (n+1)}{r (r-1)! (n-r+1) (n-r)!}$$

$$\frac{(n+1)!}{r! (n-r+1)!}$$

$$= \binom{n+1}{r}$$