

MSF Mid 1 QnA

1) What is the fermat number? find the factor of 809009 by using fermat method of factorization.

Fermat Numbers:

A number F_n is of the form

$$F_n = 2^{2^n} + 1 \quad ; \quad n \geq 0$$

is called a Fermat number.

Fermat Prime:

A Fermat number which is also a prime number is called Fermat Prime.

Examples:-

$$F_0 = 2^{2^0} + 1 = 3$$

$$F_1 = 2^{2^1} + 1 = 2^2 + 1 = 5$$

$$F_2 = 2^{2^2} + 1 = 2^4 + 1 = 17$$

$$F_3 = 2^{2^3} + 1 = 2^8 + 1 = 257$$

$$F_4 = 2^{2^4} + 1 = 2^{16} + 1 = 65537$$

$$\text{Note: } F_5 = 2^{2^5} + 1 = 2^{32} + 1 = 4294967297$$

is a composite number.

Factorise 809009 using format method of Factorization.

Given $n = 809009$.

find $\sqrt{n} = 899.45$

i) Let $t = \sqrt{n} + 1$
 $= 899 + 1$
 $t = 900$

$\therefore \Rightarrow s^2 = t^2 - n$

$s = \sqrt{991} = 31.48$

s is not a perfect square.

iii) Let $t = \sqrt{n} + 3$
 $= \sqrt{899} 899 + 3$
 $= 902$

$\Rightarrow s^2 = t^2 - n$

$s = 2\sqrt{698} = 52.8$
 $= 67.78$

$\therefore t = 904, s = 80$

~~$\therefore 903, 80$ are factors of 809009 .~~

ii) Let $t = \sqrt{n} + 2$

$\Rightarrow t = 901$

$s^2 = t^2 - n$

$s = 52.83$

vi) Let $t = \sqrt{n} + 31$
 $= 899 + 31$
 $= 903$

$s^2 = t^2 - n$

$s^2 = 903^2 - 3$

$s = \sqrt{903^2 - 3} = 80$

$n = t^2 - s^2$

$= (t+s)(t-s)$

$= 903(903+80)(903-80)$

$n = (983)(823)$

$a = 983$

$b = 823$

$\therefore 983, 823$ are the factors of 809009 .

2) Solve the system of linear eq wing chinese remainder theorems $x=2(\text{mod}3)$, $x=3(\text{mod}5)$, $x=2(\text{mod}7)$

Solve systems of congruences $x \equiv 2 \pmod{3}$ using Chinese (6)
 $x \equiv 3 \pmod{5}$ remainder Theorem.
 $x \equiv 2 \pmod{7}$

Here, $a_1 = 2, a_2 = 3, a_3 = 2$
 $n_1 = 3, n_2 = 5, n_3 = 7.$

$$n = n_1 \times n_2 \times n_3$$

$$= 3 \times 5 \times 7$$

$$n = 105$$

$$N_1 = \frac{n}{n_1} = \frac{105}{3} = 35$$

$$N_2 = \frac{n}{n_2} = \frac{105}{5} = 21$$

$$N_3 = \frac{n}{n_3} = \frac{105}{7} = 15$$

$(N_k, n_k) = 1, N_k x \equiv 1 \pmod{n_k}$ considering the
 linear congruence:

$$35x \equiv 1 \pmod{3} \quad \text{--- (1)}$$

$$21x \equiv 1 \pmod{5} \quad \text{--- (2)}$$

$$15x \equiv 1 \pmod{7} \quad \text{--- (3)}$$

① $\Rightarrow$ Let $x=1$:

$$35(1) \equiv 1 \pmod{3}$$

$$35 \equiv 1 \pmod{3}$$

$$\frac{3}{35-1} = \frac{3}{34} \times \text{not congruence.}$$

Let $x=2$: $35(2) \equiv 1 \pmod{3}$

$$70 \equiv 1 \pmod{3}$$

$$\frac{3}{70-1} = \frac{3}{69} \quad \boxed{\text{congruent}}$$

$$\therefore x_1 = 2.$$

② $\Rightarrow$ Let $x=1$: $21(1) \equiv 1 \pmod{5}$ ③ $\Rightarrow$ Let $x=1$:

$$15(1) \equiv 1 \pmod{7}$$

$$\frac{5}{21-1} = \frac{5}{20} \quad \checkmark$$

$$x_2 = 1 \quad \boxed{\text{congruent}}$$

$$\frac{7}{15-1} = \frac{7}{14} \quad \checkmark$$

$$\boxed{\text{congruent}} \quad x_3 = 1$$

Simultaneous solution of the given
System of congruence

$$\boxed{\bar{x} = a_1 n_1 x_1 + a_2 n_2 x_2 + a_3 n_3 x_3.}$$

$$= (2)(35)(2) + (3)(21)(1) + (2)(15)(1)$$

$$= 233.$$

$$x \equiv \bar{x} \pmod{n}$$

$$x \equiv 233 \pmod{105}$$

$$\frac{233}{105} = 105 \times 2 + 23$$

Let $x \Rightarrow$

$$x \equiv 23 \pmod{105}$$

$$233 \equiv 23 \pmod{105}$$

$$x = 233.$$

$$\therefore 23 \equiv 2 \pmod{3}$$

$$23 \equiv 3 \pmod{5}$$

$$23 \equiv 2 \pmod{7}$$

3) Fit a straight line $y=ax+b$ for the following data

x	0	1	2	3	4
F(x)	1	1.8	3.3	4.5	6.3

The normal equations provide an alternative method for finding the coefficients (a) and (b) in the equation ($y = ax + b$) for the least squares regression line. The normal equations are given by:

$$\sum y = a \sum x + nb$$

$$\sum xy = a \sum x^2 + b \sum x$$

values using the provided data:

$$n = 5$$

$$\Sigma x = 0 + 1 + 2 + 3 + 4 = 10$$

$$\Sigma y = 1 + 1.8 + 3.3 + 4.5 + 6.3 = 16$$

$$\Sigma xy = (0 \cdot 1) + (1 \cdot 1.8) + (2 \cdot 3.3) + (3 \cdot 4.5) + (4 \cdot 6.3) = 44.1$$

$$\Sigma x^2 = (0^2) + (1^2) + (2^2) + (3^2) + (4^2) = 30$$

Substitute these values into the normal equations:

$$16 = a \cdot 10 + 5b$$

$$44.1 = a \cdot 30 + 10b$$

Let's solve it step by step:

$$\text{Equation 1: } 16 = 10a + 5b$$

$$\text{Equation 2: } 44.1 = 30a + 10b$$

Multiply Equation 1 by 2 to make the coefficients of b match:

$$32 = 20a + 10b$$

Now, subtract Equation 2 from the modified Equation 1:

$$32 - 44.1 = (20a + 10b) - (30a + 10b)$$

Solve for a :

$$-12.1 = -10a$$

$$a = 1.21$$

Now substitute the value of a back into Equation 1 to solve for b :

$$16 = 10(1.21) + 5b$$

$$16 = 12.1 + 5b$$

$$5b = 3.9$$

$$b = 0.78$$

So, the values of a and b are 1.21 and 0.78 respectively. Therefore, the equation of the least squares regression line is

$$(y = 1.21x + 0.78).$$

4) A play on tosses 3 fair coins. He wins Rs.500 if three heads, Rs 300 if two heads appear, Rs.100 if one head occurs. on the other hand He loses Rs.1500 if 3 tails occurs. find the expected gain of player.

Let X denote the gain. Then the range $x = \left\{ \begin{matrix} -1500, \\ 100, \\ 300, \\ 500. \end{matrix} \right\}$

The sample space $S = \left\{ \begin{matrix} H H H, & T H H, \\ H T H, & T H T, \\ H T T, & T T H, \\ H H T, & T T T. \end{matrix} \right\}$

The probability of getting all three heads: (winning ₹500)

$$P(X=3) = \frac{1}{8}$$

The probability of getting 2 heads: $P(X=2) = \frac{3}{8}$ (winning ₹300)

The probability of getting 1 head $P(X=1) = \frac{3}{8}$ (winning ₹100)

The probability of getting no head (losing ₹1500)

$$P(X=0) = \frac{1}{8}$$

Required Discrete Probability Distribution

x	-1500	100	300	500
$P(x)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

$$E(x) = \sum P_i \cdot x_i$$

$$= +1500 \left(-\frac{1}{8} \right) + 100 \left(\frac{3}{8} \right) + 300 \left(\frac{3}{8} \right) + 500 \left(\frac{1}{8} \right)$$

$$= 25$$

5) find mean and variance of the uniform probability distribution given by $f(x) = 1/n$ for $x=1,2,3,...,n$

For a uniform probability distribution given by $f(x) = \frac{1}{n}$ for $x = 1, 2, 3, \dots, n$, the mean (μ) and variance (σ^2) can be calculated as follows:

1. Mean (μ):

The mean is the average value and is given by the formula:

$$\mu = \sum_{i=1}^n x_i \cdot P(x_i)$$

where x_i are the possible values of x (in this case, 1 to n) and $P(x_i)$ is the corresponding probability.

For this uniform distribution, $P(x_i) = \frac{1}{n}$ for all x_i . Therefore, the mean is the sum of all possible values divided by n :

$$\mu = \frac{1}{n} \sum_{i=1}^n x_i$$

For the given distribution:

$$\mu = \frac{1}{n} \sum_{i=1}^n i$$

$$\mu = \frac{1}{n} \sum_{i=1}^n i$$

$$\mu = \frac{1}{n} \cdot \frac{n \cdot (n+1)}{2}$$

$$\mu = \frac{n+1}{2}$$


2. Variance (σ^2):

The variance is a measure of the spread of the distribution and is given by:

$$\sigma^2 = \sum_{i=1}^n (x_i - \mu)^2 \cdot P(x_i)$$

where μ is the mean.

For this uniform distribution, $P(x_i) = \frac{1}{n}$ for all x_i . Therefore, the variance is:



Handwritten derivation of the variance formula for a uniform distribution:

$$\begin{aligned} \text{Variance} &= \sum n i x_i^2 - \mu^2 \\ &= \frac{1}{n} [1^2 + 2^2 + 3^2 + \dots + n^2] - \left(\frac{n+1}{2}\right)^2 \\ &= \frac{1}{n} \cdot \frac{n(n+1)(2n+1)}{6} - \left(\frac{n+1}{2}\right)^2 \end{aligned}$$

$$\begin{aligned}
 &= \frac{n+1}{2} \left[\frac{2n+1}{3} - \frac{n+1}{2} \right] \\
 &= \frac{n+1}{2} \left[\frac{4n+2-3n-3}{6} \right] \\
 &= \frac{(n+1)(n-1)}{12} \\
 \sigma^2 &= \frac{n^2-1}{12} \\
 \therefore \text{Mean} &= \frac{n+1}{2}, \text{ Variance} = \frac{n^2-1}{12}
 \end{aligned}$$

Therefore, the mean μ is $\frac{n+1}{2}$ and the variance σ^2 is $\frac{n^2-1}{12}$ for the given uniform probability distribution.

6) A sample of 4 items is selected at random from a box containing 12 items of which 5 are defective. Find the expected number 'E' of defective items.

Let X denote the number of defective items

among 4 items drawn from 12 items

X can take the values 0, 1, 2, 3, 4.

The number of defective items = 5

The no. of non-defective items = 7.

$$P(X=0) = P(\text{no. of defective} = 0) = \frac{{}^7C_4 {}^5C_0}{{}^{12}C_4} = \frac{35}{495} = \frac{7}{99}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

$${}^nC_1 = n$$

$${}^nC_n = 1$$

$${}^nC_0 = 1$$

$$P(X=1) = P(\text{one defect}) = \frac{{}^7C_3 {}^5C_1}{{}^{12}C_4} = \frac{175}{495} = \frac{35}{99}$$

$$P(X=2) = P(\text{two defect}) = \frac{{}^7C_2 {}^5C_2}{{}^{12}C_4} = \frac{210}{495} = \frac{14}{33}$$

$$P(X=3) = P(\text{three defect}) = \frac{{}^7C_1 {}^5C_3}{{}^{12}C_4} = \frac{70}{495} = \frac{14}{99}$$

$$P(X=4) = P(\text{all defects}) = \frac{{}^7C_0 {}^5C_4}{{}^{12}C_4} = \frac{5}{495} = \frac{1}{99}$$

Required Probability Distribution.

X	0	1	2	3	4
P(x)	$\frac{7}{99}$	$\frac{35}{99}$	$\frac{42}{99}$	$\frac{14}{99}$	$\frac{1}{99}$

$$\begin{array}{r} 1.66\overline{6} \\ 3.4\overline{8} \\ \hline 0.70\overline{70} \end{array}$$

$$\text{Mean (Expectation)} = \sum_{i=0}^4 P_i x_i$$

$$\mu = 0 + \frac{35}{99} + \frac{84}{99} + \frac{42}{99} + \frac{4}{99}$$

$$\mu = 1.\overline{6}$$

$$\text{Variance} = \sum_{i=0}^4 P_i x_i^2 - \mu^2$$

$$= 0 + \frac{35}{99} + \frac{4 \times 42}{99} + \frac{9 \times 14}{99} + \frac{16}{99} - (1.\overline{6})^2$$

$$= 3.4\overline{8} - 1.\overline{6}^2 = 0.70\overline{70}$$

7) What is the fermat number? find the factor of 426749 by using formal method of factorization.

Fermat Numbers:

A number F_n is of the form

$$F_n = 2^{2^n} + 1 \quad ; \quad n \geq 0$$

is called a Fermat number.

Fermat Prime:

A Fermat number which is also a prime number is called Fermat Prime.

Examples:-

$$F_0 = 2^{2^0} + 1 = 3$$

$$F_1 = 2^{2^1} + 1 = 2^2 + 1 = 5$$

$$F_2 = 2^{2^2} + 1 = 2^4 + 1 = 17$$

$$F_3 = 2^{2^3} + 1 = 2^8 + 1 = 257$$

$$F_4 = 2^{2^4} + 1 = 2^{16} + 1 = 65537$$

$$\text{Note: } F_5 = 2^{2^5} + 1 = 2^{32} + 1 = 4294967297$$

is a composite number.

factor of 426749 by using formal method of factorization:

1. Choose t :

Let's choose $t = 655$ (you can experiment with different values).

2. Check $t^2 - N$:

Calculate $t^2 - N$ and check if it's a perfect square. If

$t^2 - N = s^2$ for some s , then N can be factored as

$(t + s)(t - s)$.

$$656^2 - 426749 = 430336 - 426749 = 3587$$

Since 3587 is not a perfect square, we need to choose a different value for t and repeat the process.

3. Choose a Different t :

Let's try another value. Let's choose $t = 657$.

4. Check $t^2 - N$:

$$657^2 - 426749 = 431649 - 426749 = 4900$$

Now, $t^2 - N = 4900$, which is a perfect square (70^2).

5. Factorize Using the Result:

Since $s^2 = t^2 - N = 4900$ and $4900 = 70^2$, we can write N as the difference of two squares:

$$N = t^2 - s^2 = (t + 70)(t - 70)$$

Therefore, the factors of 426749 are $t + 70$ and $t - 70$.

6. Calculate Factors:

$$a + 70 = 657 + 70 = 727$$

$$a - 70 = 657 - 70 = 587$$

Therefore, the factors of 426749 are 587 and 727.

8) solve the system of linear equation using chinese remainder theorem $x \equiv 5 \pmod{6}$, $x \equiv 1 \pmod{11}$ $x \equiv 3 \pmod{17}$

$$\begin{aligned} \text{a) } x &\equiv 5 \pmod{6} & \text{comparing to } x &\equiv a_1 \pmod{n_1} \\ x &\equiv 4 \pmod{11} & x &\equiv a_2 \pmod{n_2} \\ x &\equiv 3 \pmod{17} & x &\equiv a_3 \pmod{n_3} \end{aligned}$$

$$a_1 = 5, a_2 = 4, a_3 = 3$$

$$n_1 = 6, n_2 = 11, n_3 = 17$$

$$N = n_1 \times n_2 \times n_3$$

$$N = 6 \times 11 \times 17$$

$$N = 1122$$

$$N_1 = \frac{N}{n_1} \Rightarrow N_1 = \frac{1122}{6} = 187$$

$$N_2 = \frac{N}{n_2} \Rightarrow N_2 = \frac{1122}{11} = 102$$

$$N_3 = \frac{N}{n_3} \Rightarrow N_3 = \frac{1122}{17} = 66$$

$$\text{Here, } (N_k, n_k) = 1 \text{ i.e., } N_k \equiv 1 \pmod{n_k}$$

$\Rightarrow$

$$187 x_1 \equiv 1 \pmod{6} \quad \text{--- (1)}$$

$$102 x_2 \equiv 1 \pmod{11} \quad \text{--- (2)}$$

$$66 x_3 \equiv 1 \pmod{17} \quad \text{--- (3)}$$

For Eq (1)

For Eq (2)

$$\text{i) If } x_1 = 1: 187(1) \equiv 1 \pmod{6} \quad \text{ii) If } x_2 = 1: 102(1) \equiv 1 \pmod{11}$$

$$\Rightarrow 187 \pmod{6} = 1$$

$$\Rightarrow 102 \pmod{11} = 1$$

$$1 = 1$$

$$3 \neq 1$$

not congruent

It is congruent

ii)

$$\text{If } x_2 = 1: 102(1) \equiv 1 \pmod{11}$$

$$x_1 = 1$$

$$102 \cdot 2 = 204 \equiv 1 \pmod{11}$$

$$204 \pmod{11} = 1$$

$$6 \neq 1$$

not congruent

$$\text{iii) If } x_2 = 3 : 102(3) \equiv 1 \pmod{11}$$

$$306 \pmod{11} = 1$$

$$9 \neq 1$$

It is not congruent

$$\text{iv) If } x_2 = 4 : 102(4) \equiv 1 \pmod{11}$$

$$408 \pmod{11} = 1$$

$$1 = 1$$

It is congruent

$$x_2 = 4.$$

For Eq (3)

$$\text{i) If } x_3 = 1 : 66(1) \equiv 1 \pmod{17}$$

$$66 \pmod{17} = 1$$

$$15 \neq 1$$

not congruent

$$\text{ii) If } x_3 = 2 : 66(2) \equiv 1 \pmod{17}$$

$$132 \pmod{17} = 1$$

$$13 \neq 1$$

not congruent

$$\text{iii) If } x_3 = 7 : 66(7) \equiv 1 \pmod{17}$$

$$462 \pmod{17} = 1$$

$$3 \neq 1$$

not congruent

$$\text{iv) If } x_3 = 8 : 66(8) \equiv 1 \pmod{17}$$

$$528 \pmod{17} = 1$$

$$1 = 1$$

If is congruent

$$x_3 = 8$$

Hence,

$$x_1 = 1$$

$$x_2 = 4$$

$$x_3 = 8$$

~~and~~ $\begin{matrix} 187 & 102 & 66 \\ a_1 = 5, a_2 = 4, a_3 = 3 \\ n_1 = 6, n_2 = 11, n_3 = 17 \end{matrix}$

according to simultaneous solution of the given system of congruence.

$$\bar{x} = a_1 N_1 x_1 + a_2 N_2 x_2 + a_3 N_3 x_3$$

$$x = \bar{x} \pmod{N} \quad (\text{or}) \quad \bar{x} \equiv x \pmod{N}$$

$$a_1 = 5, a_2 = 4, a_3 = 3$$

$$N_1 = 187, N_2 = 102, N_3 = 66$$

$$x_1 = 1, x_2 = 4, x_3 = 8$$

$$\bar{x} = 5(187)(1) + 4(102)(4) + 3(66)(8)$$

$$\bar{x} = 935 + 1632 + 1584$$

$$\bar{x} = 4151$$

$$\therefore N = ~~1122~~ 1122$$

$$4151 \equiv x \pmod{1122}$$

$$x = 4151 \pmod{1122}$$

$$x = 785$$

Therefore, The system of linear equation is

$$785 \equiv 5 \pmod{6}$$

$$785 \equiv 4 \pmod{11}$$

$$785 \equiv 3 \pmod{17}$$

9) Out of 800 families with 5 children each how would many you expect to have a) 3 boys b) 5 girls c) either 2 (or) 3 boys d) atleast one boy? Assume equal properties probabilities for boys and girls.

Let probability of no. of boys in each family be $p = \frac{1}{2}$, of girls $q = \frac{1}{2}$.

no. of boys in family = X .

no. of children $= n = 5$.

i) Probability of having 3 boys:

$$P(X=3)$$

$$P(X=3) = {}^5C_3 \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^{5-3}$$

$$= \frac{5}{16}$$

For 800 families, the probability of no. of families having 3 boys:

$$\Rightarrow \frac{5}{16} \times 800 = 250 \text{ families.}$$

ii) 5 girls

$$P(X=0) = {}^5C_0 \left(\frac{1}{2}\right)^0 \left(\frac{1}{2}\right)^{5-0}$$

$$= \frac{1}{32}$$

For 800 families, the no. of families having 5 girls

$$\text{i.e., } \frac{1}{32} \times 800 = 25$$

iii) 2 or 3 boys

$$= P(X=2) + P(X=3)$$

$$= {}^5C_2 \left(\frac{1}{2}\right)^2 \left(\frac{1}{2}\right)^{5-2} + {}^5C_3 \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^{5-3}$$

$$= \frac{5}{16} + \frac{5}{16}$$

$$= \frac{10}{16} = \frac{5}{8}$$

For 800 families, the probability of no. of families having 2 or 3 boys is $\frac{10}{16} \times 800$

iv) Atleast one boy

$$P(X \geq 1) = 1 - P(X=0)$$

$$= 1 - \frac{1}{32}$$

$$= \frac{31}{32}$$

For 800 families the probability of no. of families having atleast one boy is

$$\frac{31}{32} \times 800 = 775.$$

10) Write the proof mean of normal distribution.

Mean of the Normal Distribution:-

By the definition of Normal distribution,

$$f(x) = \frac{1}{\sigma \sqrt{2\pi}} e^{-\frac{1}{2} \left(\frac{x-\mu}{\sigma} \right)^2}, \quad -\infty < x < \infty, \quad -\infty < \mu < \infty$$

$$\text{Mean } \mu = E(x) = \int_{-\infty}^{\infty} x f(x) dx$$

$$= \int_{-\infty}^{\infty} x \frac{1}{\sigma \sqrt{2\pi}} e^{-\frac{1}{2} \left(\frac{x-\mu}{\sigma} \right)^2} dx$$

$$= \frac{1}{\sigma \sqrt{2\pi}} \int_{-\infty}^{\infty} x e^{-\frac{1}{2} \left(\frac{x-\mu}{\sigma} \right)^2} dx$$

$$\begin{aligned} \text{Let } \frac{x-\mu}{\sigma} &= z \Rightarrow x-\mu = z\sigma \\ x &= z\sigma + \mu \\ dx &= \sigma dz + 0. \end{aligned}$$

$$\Rightarrow \frac{1}{\cancel{\sigma} \sqrt{2\pi}} \int_{-\infty}^{\infty} (\cancel{\sigma} z + \mu) e^{-\frac{1}{2} z^2} (\cancel{\sigma} dz)$$

$$= \frac{\sigma}{\sqrt{2\pi}} \int_{-\infty}^{\infty} z e^{-\frac{z^2}{2}} dz + \frac{\mu}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{z^2}{2}} dz$$

$$= 0 + \frac{2\mu}{\sqrt{2\pi}} \int_0^{\infty} e^{-\frac{z^2}{2}} dz$$

$$\begin{aligned} \text{Let } \frac{z^2}{2} &= t \Rightarrow z = \sqrt{2t} \\ 2z dz &= 2 \cdot dt \Rightarrow dz = \frac{dt}{z} \\ dz &= \frac{dt}{\sqrt{2t}} \end{aligned}$$

$$= \frac{2\mu}{\sqrt{2\pi}} \int_0^{\infty} e^{-t} \frac{dt}{\sqrt{2t}} \quad \left[\because \int_0^{\infty} e^{-x} x^{n-1} dx = \frac{\Gamma(n)}{n} \right]$$

$$\frac{2\mu}{\sqrt{2} \sqrt{2\pi}} \int_0^{\infty} e^{-t} t^{-1/2} dt$$

$$\frac{\mu}{\sqrt{\pi}} \int_0^{\infty} e^{-t} t^{1/2-1} dt$$

$$\begin{aligned} n-1 &= -\frac{1}{2} \\ n &= \frac{1}{2} \\ n > \frac{1}{2} \end{aligned}$$

$$= \frac{\mu}{\sqrt{\pi}} \sqrt{\frac{1}{2}} \quad \left[\because \int_0^{\infty} e^{-t} t^{n-1} dt = \frac{\Gamma(n)}{n} \right]$$

$$= \frac{\mu}{\sqrt{\pi}} \sqrt{\pi}$$

$$\text{Mean} = \mu$$

$$\begin{aligned} \int_0^{\infty} e^{-t} t^{n-1} dt &= \frac{\Gamma(n)}{n} \\ \int_0^{\infty} e^{-t} t^{1/2-1} dt &= \frac{\Gamma(1/2)}{1/2} \\ \int_0^{\infty} e^{-t} t^{-1/2} dt &= \frac{\Gamma(1/2)}{1/2} \end{aligned}$$

11) Write proof variance of normal distribution

Variance of Normal Distribution:

By the definition of Normal Distribution:

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}, \quad -\infty < x < \infty$$

$-\infty < \mu < \infty$

$$\text{Variance } \sigma^2 = \int_{-\infty}^{\infty} (x-\mu)^2 f(x) dx$$

$$= \int_{-\infty}^{\infty} (x-\mu)^2 \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} dx$$

$$= \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} (x-\mu)^2 e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} dx$$

Put $\frac{x-\mu}{\sigma} = z \Rightarrow$

$$x = \sigma z + \mu$$
$$dx = \sigma dz$$

$$= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (\sigma z)^2 e^{-\frac{1}{2}(z)^2} dz$$

$$= \frac{\sigma^2}{\sqrt{2\pi}} \int_{-\infty}^{\infty} z^2 e^{-\frac{z^2}{2}} dz$$

Put $\frac{z^2}{2} = t$

$$z = \sqrt{2t}$$

$$dz = \frac{dt}{\sqrt{2t}}$$

$$= \frac{2\sigma^2}{\sqrt{2\pi}} \int_0^{\infty} z^2 e^{-z^2/2} dz$$

$$= \frac{2\sigma^2}{\sqrt{2\pi}} \int_0^{\infty} (\sqrt{2t})^2 e^{-t} \frac{dt}{\sqrt{2t}}$$

$$= \frac{2 \times 2 \times \sigma^2}{\sqrt{2} \sqrt{2} \pi} \int_0^{\infty} e^{-t} \frac{t}{t^{1/2}} dt$$

$$= \frac{2\sigma^2}{\sqrt{\pi}} \int_0^{\infty} e^{-t} t^{3/2-1} dt$$

$$= \frac{2\sigma^2}{\sqrt{\pi}} \sqrt{\frac{3}{2}}$$

$$\left[\because \sqrt{\frac{3}{2}} = \frac{1}{2} \sqrt{\frac{1}{2}} = \frac{1}{2} \sqrt{\pi} \right]$$

$$= \frac{2\sigma^2}{\sqrt{\pi}} \cdot \frac{1}{2} \sqrt{\pi}$$

$$n = -1 + \frac{1}{2}$$

$$n = \frac{1}{2} + 1$$

$$n = 3/2$$

$$\sqrt{n} = (n-1) \sqrt{n-1}$$

$$\text{Variance} = \sigma^2$$

$$\sqrt{\frac{1}{2}} = \sqrt{\pi}$$

calculate expectation and variance of 'X' variance of the probability distribution of the random variable 'X' is given by

x	-1	0	1	2	3
F(x)	0.3	0.1	0.1	0.3	0.2

Given the probability distribution of X:

x	-1	0	1	2	3
F(x)	0.3	0.1	0.1	0.3	0.2

1. Expectation (μ):

$$\mu = E(X) = \sum_i x_i \cdot P(X = x_i)$$

$$\mu = (-1 \cdot 0.3) + (0 \cdot 0.1) + (1 \cdot 0.1) + (2 \cdot 0.3) + (3 \cdot 0.2)$$

$$\mu = -0.3 + 0 + 0.1 + 0.6 + 0.6$$

$$\mu = 1$$

2. Variance (σ^2):

$$\sigma^2 = \sum_i x_i^2 \cdot P(X = x_i) - \mu^2$$

Given that $\mu = 1$ (calculated previously), let's use this formula to find the variance:

$$\sigma^2 = \sum_i x_i^2 \cdot P(X = x_i) - \mu^2$$

$$\sigma^2 = (-1)^2 \cdot 0.3 + (0)^2 \cdot 0.1 + (1)^2 \cdot 0.1 + (2)^2 \cdot 0.3 + (3)^2 \cdot 0.2 - 1^2$$

$$\sigma^2 = 0.3 + 0 + 0.1 + 1.8 + 1.2 - 1$$

$$\sigma^2 = 2.4$$

So, the expectation (μ) is 1, and the variance (σ^2) is 2.4 for the given probability distribution.

12) A sample of u items is selected a vandom from a box containing 12 items of which 5 are defective. Find

the expected number 'E' of defective items

// Same Question 6

13) In normal distribution define mean =
median = mode

// Not given